

OPT

Olopatadine 0.7%

Sterile & Pyrogen Free Eye Drops

Composition

OPT Sterile Eye Drops: Each mL contains Olopatadine Hydrochloride USP equivalent to Olopatadine 7 mg.

Preservative: Sodium Perborate 0.01%

Pharmacology

Olopatadine is an antihistamine with mast cell stabilizing properties. It inhibits histamine and prevents histamine induced inflammatory cytokine production.

Indications

Ocular itching associated with allergic conjunctivitis.

Dosage and Administration

One drop in the affected eye(s) once daily or as advised by the registered physician.



Wash hands before using the eye drops



Pull down your eyelid with a clean finger



Bring the dropper tip close to the eye



Gently press the bottom of the dropper to release of medicine



Gently close your eyes 2-3 minutes after administration

Contraindications

Contraindicated in patients who are hypersensitive to this drug or to any ingredients in the formulation.

Side Effects

The following adverse effect has been reported:

Headache, stinging or burning of eye and somnolence.

Warnings & Precautions

To prevent contaminating the dropper tip and solution, care should be taken not to touch the eyelids or surrounding areas with the dropper tip of the bottle. Patients should be advised not to wear a contact lens if their eye is red.



Do not pierce with needles



Do not cut with scissor or knife



Do not touch the nozzle/dropper tip



Close the upper part of the cap with the container just after the use



After 30 days using please keep the container in bin

Special Guideline

- ❖ If more than one topical ophthalmic product is to be used, other eye drops should be instilled at least 5 to 10 minutes apart.
- ❖ If using two types of eye medications such as solution and suspension, use the solution first and then use suspension.
- ❖ If using eye drops and an eye ointment, use the eye drops first then apply the ointment at least 10 minutes later.
- ❖ Don't share eye drops with another persons because sharing drops could spread germs and infection.

Use in Pregnancy & Lactation

Pregnancy: There are no adequate or well-controlled studies with olopatadine in pregnant women. Should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Lactation: Caution should be exercised when olopatadine is administered to a lactating mother.

Use in Children & Adolescents

Safety and effectiveness in pediatric patients below the age of 2 years have not been established.

Drug Interactions

No Drug interaction studies have been performed.

Overdose

No information is available.

Storage

Store between 4°-25°C, away from light and in a dry place. Keep out of the reach of children.

Packaging

OPT Sterile Eye Drops: Each LDPE dropper bottle contains 5 mL Sterile & Pyrogen free Eye Drops and a patient information leaflet.

Manufactured by:

UNIDO
Pharmaceutical Limited
Batpara, Narsingdi, Bangladesh.

PM 2009

