

Uniprost

Travoprost USP 0.004%

Sterile & Pyrogen Free Eye Drops

Composition

Uniprost sterile eye drops: Each mL contains Travoprost USP 0.04 mg.

Preservative: Ionic Buffered System

Pharmacology


Travoprost, an isopropyl ester prodrug, is rapidly hydrolyzed by esterases in the cornea to the biologically active free acid. Travoprost free acid is a highly selective, potent agonist for the FP prostanoid receptor. FP receptor agonists are thought to reduce intraocular pressure (IOP) by increasing the outflow of aqueous humor, primarily by increased uveoscleral outflow.

Indications

Reduction of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma.

Doses & Administration

One drop in the affected eye(s) once daily preferably in the evening or as advised by the registered physician.



Contraindications

Travoprost is contraindicated in patients with hypersensitive to any excipients of this preparation.

Side Effects

The most common ocular adverse effect is brown pigmentation of the iris, abnormal eyelash growth, prostaglandin associated periorbitopathy (PAP) syndrome, conjunctival hyperemia, skin hyperpigmentation, dry eye, eye discomfort and eye disorders.

Warnings & Precautions

- Travoprost should be used with caution in patients with active intraocular inflammation such as Uveitis.
- Travoprost should be used with caution in patients with known risk factors for macular edema.
- Contact lenses should be removed prior to installation of Travoprost and may be reinserted 15 minutes following its administration.



Special Guideline

- ❖ If more than one topical ophthalmic product is to be used, other eye drops should be instilled at least 5 to 10 minutes apart.
- ❖ If using two types of eye medications such as solution and suspension, use the solution first and then use suspension.
- ❖ If using eye drops and an eye ointment, use the eye drops first then apply the ointment at least 10 minutes later.
- ❖ Don't share eye drops with another persons because sharing drops could spread germs and infection.

Use in Pregnancy & Lactation

Pregnancy: There are no adequate and well-controlled clinical study in pregnant women. Travoprost should be administered during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Lactating: It is not known whether the drug or its metabolites are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Travoprost is administered to a lactating women.

Use in Children & Adolescent

The efficacy and safety of Travoprost eye drops in patients below the age of 16 years have not been established.

Drug Interaction

No specific interaction was observed with **Uniprost** eye drops.

Overdose

No cases of overdose have been reported.

Storage

Store at 2°C - 25°C away from light and in a dry place. Keep out of the reach of children.

Packaging

Uniprost sterile eye drops: Each LDPE dropper bottle contains 3 mL Sterile & Pyrogen free Eye Drops and a patient information leaflet.

Manufactured by:

UNIDO
Pharmaceutical Limited
Batpara, Narsingdi, Bangladesh.

PM 2027

