

# Vigalon

Moxifloxacin 0.5%

## Sterile & Pyrogen Free Eye Drops

### Composition

**Vigalon Sterile Eye Drops:** Each mL drops contains Moxifloxacin Hydrochloride USP equivalent to Moxifloxacin 5 mg.

**Preservative:** Product is self preserved

### Pharmacology


Moxifloxacin active in vitro against a broad spectrum of Gram-positive and Gram-negative ocular pathogens, atypical microorganisms and anaerobes. It works against topoisomerase II and topoisomerase IV.

### Indications

Bacterial conjunctivitis, Blepharitis, Dacryocystitis, Hordeolum (Stye), Tarsadenitis, Keratitis as well as preoperative & postoperative prophylaxis.

### Dosage & Administration

**Vigalon Eye Drops:** One drop in the affected eyes, 3 times a day for 7 days or as advised by the registered physician.



### Contraindications

**Moxifloxacin Eye Drops** is contraindicated in patients with a history of hypersensitivity to Moxifloxacin, to other quinolones or to any of the components in this medication.

### Side Effects

After ocular use the following adverse effect has been reported: Burning or discomfort, keratopathy, corneal staining, corneal precipitates or infiltrates and photophobia.

### Warnings & Precautions

As with other anti-infectives, prolonged use may result in overgrowth of non-susceptible organisms, including fungi.



### Special Guideline

- ❖ If more than one topical ophthalmic product is to be used, other eye drops should be instilled at least 5 to 10 minutes apart.
- ❖ If using two types of eye medications such as solution and suspension, use the solution first and then use suspension.
- ❖ If using eye drops and an eye ointment, use the eye drops first then apply the ointment at least 10 minutes later.
- ❖ Don't share eye drops with another persons because sharing drops could spread germs and infection.

### Use in Pregnancy & Lactation

**Pregnancy:** **Moxifloxacin Eye Drops** should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus as there are no adequate and well-documented studies in pregnant women.

**Lactation:** Moxifloxacin has not been measured in human milk, although it can be presumed to be excreted in human milk. Caution should be exercised when **Moxifloxacin Eye Drops** are administered to a lactating mother.

### Use in Children & Adolescents

**Moxifloxacin Eye Drops:** Safe and well tolerated in pediatric patients, including all age categories: newborns (0-27 days), infants and toddlers (28 days-23 months), children (2-11 years) and adolescents (12-17 years).

### Drug Interactions

Drug-drug interaction studies have not been conducted with **Moxifloxacin Eye Drops**

### Overdose

No information is available.

### Storage

Store below 30°C temperature, away from light and in a dry place. Keep out of the reach of children.

### Packaging

**Vigalon Sterile Eye Drops:** Each LDPE dropper bottle contains 5 mL Sterile & Pyrogen free Eye Drops and a patient information leaflet.

Manufactured by:

**UNIDO**  
Pharmaceutical Limited  
Batpara, Narsingdi, Bangladesh.

PM 2003

